

HOW TO TAKE DOPTELET

Your guide to taking DOPTELET before your procedure



What is DOPTELET® (avatrombopag) [dop-TELL-it]?

DOPTELET is a prescription medicine used to treat low blood platelet counts in adults with long-lasting (chronic) liver disease who are scheduled to have a medical or dental procedure.

DOPTELET is not used to make platelet counts normal in adults with chronic liver disease. It is not known if DOPTELET is safe and effective in children.

IMPORTANT SAFETY INFORMATION FOR DOPTELET

Before you take DOPTELET, tell your healthcare provider about all of your medical conditions, including if you:

- have ever had a blood clot.
- are pregnant or plan to become pregnant. DOPTELET may harm your unborn baby. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with DOPTELET.
- are breastfeeding or plan to breastfeed. It is not known if DOPTELET passes into your breast milk. Do not breastfeed during your treatment with DOPTELET and for at least 2 weeks after the last dose. Talk to your healthcare provider about the best way to feed your baby during this time.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. DOPTELET may affect the way other medicines work, and other medicines may affect the way DOPTELET works.

Please see rest of Important Safety Information on page 4.
Full Prescribing Information for DOPTELET® (avatrombopag) is available at DOPTELET.com.

How should I take DOPTELET?

You should take DOPTELET® (avatrombopag) just as your healthcare provider prescribed.

Three things you should know about taking DOPTELET:



ONCE A DAY.

Take your dose of DOPTELET once a day.



FOR 5 DAYS.

Take DOPTELET for 5 days in a row as directed by your healthcare provider.



WITH FOOD.

Take DOPTELET every day with food.

- If you are taking DOPTELET prior to a scheduled medical procedure and you miss a dose, contact your healthcare provider for further dosing instructions.
- Complete all 5 days of medication.

Starting on the right day is important.

Someone from your healthcare team, most likely the specialty pharmacy, will contact you to remind you when to start taking DOPTELET.

If you have any questions about how or when to take DOPTELET, call your healthcare provider. You can also call your specialty pharmacy, or Doptelet CONNECT™ at 1-833-368-2663 and they will help you.

IMPORTANT SAFETY INFORMATION FOR DOPTELET (CONT.)

How should I take DOPTELET?

- Take DOPTELET exactly as your healthcare provider tells you to take it.
- Your healthcare provider will tell you how much DOPTELET to take and when to start taking it.
- Your healthcare provider may change your dose of DOPTELET depending on your blood platelet counts.
- Take DOPTELET with food.
- If you take DOPTELET to treat your low blood platelet counts due to chronic liver disease before a medical or dental procedure, your healthcare provider will check your platelet count before treatment and on the day of your scheduled procedure.
- If you are taking DOPTELET prior to a scheduled medical procedure and you miss a dose, contact your healthcare provider for further dosing instructions.
- If you take too much DOPTELET, call your healthcare provider or go to the nearest hospital emergency room right away.

**Please see rest of Important Safety Information on page 4.
Full Prescribing Information for DOPTELET® (avatrombopag) is available at DOPTELET.com.**

DOPTELET 40 mg
2 tablets a day for 5 days

DOPTELET 60 mg
3 tablets a day for 5 days



Your DOPTELET schedule will look like this.

Speak to your healthcare provider about the exact day you should start DOPTELET. Your specialty pharmacy will contact you to remind you of those days. You can use the schedule below to write in your days. This may help you remember.

DAY 1:

DAY 2:

DAY 3:

DAY 4:

DAY 5:

PROCEDURE DAY:

Your healthcare provider may change your dose of DOPTELET, depending on your blood platelet counts.

IMPORTANT SAFETY INFORMATION FOR DOPTELET (CONT.)

What are the possible side effects of DOPTELET?

DOPTELET may cause serious side effects, including blood clots. People with chronic liver disease and people with certain blood clotting conditions may have an increased risk of developing blood clots. Tell your healthcare provider right away if you have signs and symptoms of a blood clot, including:

- swelling, pain, or tenderness in your legs
- fast heartbeat
- shortness of breath
- stomach (abdominal) pain or tenderness
- chest pain

The most common side effects of DOPTELET in people with chronic liver disease are:

- fever
- headache
- stomach (abdominal) pain
- tiredness
- nausea
- swelling of the hands or feet

These are not all the possible side effects of DOPTELET.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see rest of Important Safety Information on page 4. Full Prescribing Information for DOPTELET® (avatrombopag) is available at DOPTELET.com.

IMPORTANT SAFETY INFORMATION FOR DOPTELET (CONT.)

How should I store DOPTELET?

- Store DOPTELET at room temperature between 68°F to 77°F (20°C to 25°C).
- Store DOPTELET tablets in the original package.

Keep DOPTELET and all medicines out of the reach of children.

General information about the safe and effective use of DOPTELET.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use DOPTELET for a condition for which it was not prescribed. Do not give DOPTELET to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about DOPTELET that is written for health professionals.

What are the ingredients in DOPTELET?

Active ingredient: avatrombopag

Inactive ingredients: lactose monohydrate, colloidal silicon dioxide, crospovidone, magnesium stearate and microcrystalline cellulose. Tablet coating film: polyvinyl alcohol, talc, polyethylene glycol, titanium dioxide and ferric oxide yellow.

Please see full [Prescribing Information](#), including the [Patient Information leaflet](#).

This Patient Information has been approved by the U.S. Food and Drug Administration.

For product information and adverse event reports, please contact:

Sobi Medical Information Call Center

Phone: 1-866-773-5274

E-mail: medinfo.us@sobi.com

Reference: DOPTELET [package insert]. Durham, NC: AkaRx, Inc.

If you have any questions about how or when to take DOPTELET, call your healthcare provider. You can also call your specialty pharmacy, or Doptelet Connect at 1-833-368-2663 and they will help you.



Doptelet is a registered trademark of AkaRx, Inc.
Sobi is a trademark of Swedish Orphan Biovitrum AB (publ)
© 2021 Sobi, Inc. – All rights reserved

PP-13744 12/21 Printed in the USA